

**WELCOME**  
**TO**



**Pharmacy**

Learn and Educate



## Bachelor of Pharmacy Human Anatomy and Physiology II

### NOTES

- ✓ Unit 1
  - ✓ Unit 2
  - ✓ Unit 3
  - ✓ Unit 4
  - ✓ Unit 5
- All Unit  
in  
One PDF**

Visit our Website

[WWW.fdpharmacy.in](http://WWW.fdpharmacy.in)



## Bachelor of Pharmacy Pathophysiology

### NOTES

- ✓ Unit 1
  - ✓ Unit 2
  - ✓ Unit 3
  - ✓ Unit 4
  - ✓ Unit 5
- All Unit  
in  
One PDF**

Visit our Website

[WWW.fdpharmacy.in](http://WWW.fdpharmacy.in)



## Bachelor of Pharmacy Pharmaceutical Organic Chemistry I

### NOTES

- ✓ Unit 1
  - ✓ Unit 2
  - ✓ Unit 3
  - ✓ Unit 4
  - ✓ Unit 5
- All Unit  
in  
One PDF**

Visit our Website

[WWW.fdpharmacy.in](http://WWW.fdpharmacy.in)



## Bachelor of Pharmacy Biochemistry

### NOTES

- ✓ Unit 1
  - ✓ Unit 2
  - ✓ Unit 3
  - ✓ Unit 4
  - ✓ Unit 5
- All Unit  
in  
One PDF**

Visit our Website

[WWW.fdpharmacy.in](http://WWW.fdpharmacy.in)



## Bachelor of Pharmacy Computer Applications in Pharmacy

### NOTES

- ✓ Unit 1
  - ✓ Unit 2
  - ✓ Unit 3
  - ✓ Unit 4
  - ✓ Unit 5
- All Unit  
in  
One PDF**

Visit our Website

[WWW.fdpharmacy.in](http://WWW.fdpharmacy.in)



## Bachelor of Pharmacy Environmental Sciences

### NOTES

- ✓ Unit 1
  - ✓ Unit 2
  - ✓ Unit 3
  - ✓ Unit 4
  - ✓ Unit 5
- All Unit  
in  
One PDF**

Visit our Website

[WWW.fdpharmacy.in](http://WWW.fdpharmacy.in)





# FD Pharmacy

.....

## D.Pharma B.Pharma



- 👉 PDF Notes
- 👉 Practical Manual
- 👉 Important Questions
- 👉 Assignment etc

 Download Now



[www.fdp pharmacy.in](http://www.fdp pharmacy.in)

# BIOCHEMISTRY


## UNIT 2

TOPIC :

- **Biological oxidation**

Electron transport chain (ETC) and its mechanism.

level Oxidative phosphorylation & its mechanism and substrate phosphorylation Inhibitors ETC and oxidative phosphorylation/Uncouplers



Pharmacy  
Learn and Educate

## Biological Oxidation

- Biological oxidation is the process by which cells release energy from nutrients (mainly carbohydrates, fats, and proteins) via redox reactions.
- It involves oxidation of substrates to produce ATP, the energy currency of cells.
- The final stage of this process is the Electron Transport Chain (ETC) in the mitochondria.
- Oxidation = loss of electrons or hydrogen
- Reduction = gain of electrons or hydrogen
- Redox reactions drive the flow of electrons, releasing energy
- Coenzymes like  $\text{NAD}^+$  and  $\text{FAD}$  act as electron carriers
  - $\text{NAD}^+ \rightarrow \text{NADH}$
  - $\text{FAD} \rightarrow \text{FADH}_2$

## Electron Transport Chain (ETC)

- The Electron Transport Chain (ETC) is the final step of aerobic respiration.
- It is also known as the Electron Transport System (ETS).
- Occurs in the inner mitochondrial membrane of eukaryotic cells.
- It is a series of protein complexes and mobile electron carriers.
- Electrons are transferred from molecules with low redox potential to those with high redox potential.
- This transfer of electrons releases energy, which is used to pump protons ( $\text{H}^+$ ) across the membrane.
- The energy stored in the resulting proton gradient is used to generate ATP by a process called oxidative phosphorylation.

## Main Complexes of ETC

### Complex I – NADH: Ubiquinone Oxidoreductase

- Accepts electrons from **NADH**
- Transfers electrons to **Coenzyme Q (Ubiquinone)**
- **Pumps 4 H<sup>+</sup> ions** from the mitochondrial matrix to the intermembrane space
- Also called **NADH dehydrogenase**

### Complex II – Succinate Dehydrogenase

- Accepts electrons from **FADH<sub>2</sub>** (produced in the TCA cycle)
- Transfers electrons to **Coenzyme Q**
- **Does NOT pump protons**
- Also part of the **TCA cycle**

### Complex III – Cytochrome bc<sub>1</sub> Complex

- Accepts electrons from **Coenzyme Q**
- Transfers electrons to **Cytochrome c**
- **Pumps 4 H<sup>+</sup> ions** into the intermembrane space

### Complex IV – Cytochrome c Oxidase

- Accepts electrons from **Cytochrome c**
- Transfers electrons to **molecular oxygen (O<sub>2</sub>)** → **Forms water (H<sub>2</sub>O)**
- **Pumps 2 H<sup>+</sup> ions**
- Final complex in the chain

## Mechanism of ETC

- ❖ NADH donates electrons to Complex I → electrons move to Coenzyme Q, 4 H<sup>+</sup> are pumped into intermembrane space.
- ❖ FADH<sub>2</sub> donates electrons to Complex II → passes them to Coenzyme Q (no H<sup>+</sup> pumping here).
- ❖ Coenzyme Q transfers electrons to Complex III → 4 H<sup>+</sup> are pumped into intermembrane space.
- ❖ Electrons are carried by Cytochrome c to Complex IV → 2 H<sup>+</sup> are pumped, and O<sub>2</sub> is reduced to water.
- ❖ Accumulated H<sup>+</sup> ions in intermembrane space generate proton motive force (PMF).

## Significance of ETC and Biological Oxidation

- Major source of **ATP** production in aerobic organisms
- Essential for:
  - Muscle contraction
  - Nerve impulse conduction
  - Biosynthesis of macromolecules
- ETC is **aerobic** – requires **oxygen** as final electron acceptor
- Generates **heat** (thermogenesis) in brown fat (non-shivering)

Learn and Educate

## Oxidative Phosphorylation

- Oxidative phosphorylation is the process of ATP synthesis that occurs at the end of the Electron Transport Chain (ETC).
- It uses the energy from electrons transferred through the ETC to pump protons ( $H^+$ ) across the inner mitochondrial membrane.
- The resulting proton gradient powers ATP synthase to form ATP from  $ADP + P_i$ .
- It is called “oxidative” because it requires oxygen as the final electron acceptor.

### Mechanism of Oxidative Phosphorylation

The process involves two major components:

#### A. Electron Transport Chain (ETC)

- NADH and  $FADH_2$  donate electrons to the ETC.
- As electrons move through Complexes I, III, and IV:
  - **Protons ( $H^+$ )** are pumped from the mitochondrial matrix into the **intermembrane space**.
  - Creates a **proton gradient (proton motive force)**.

#### B. Chemiosmosis and ATP Synthesis

- The protons flow back into the matrix through **ATP synthase (Complex V)**.
- This flow drives the formation of ATP:



#### C. Final Step

- At **Complex IV**, electrons combine with  $O_2$  and protons to form  $H_2O$ .
- **Oxygen is the final electron acceptor.**

## Significance of Oxidative Phosphorylation

- Produces most of the ATP in aerobic respiration ( $\approx 26-28$  ATP per glucose).
- Essential for energy-demanding processes: muscle contraction, nerve transmission, biosynthesis.
- Maintains cellular energy balance and life-sustaining functions.



Pharmacy

Learn and Educate

## Substrate-Level Phosphorylation (SLP)

**Substrate-Level Phosphorylation (SLP)** is the direct synthesis of ATP (or GTP) by the transfer of a phosphate group from a high-energy phosphorylated intermediate to ADP (or GDP).


- Unlike oxidative phosphorylation, **SLP does not require oxygen** or an electron transport chain.
- The phosphate group is donated **directly by the substrate molecule** via enzyme-catalyzed reactions.

### Examples of SLP Reactions

#### 1. Glycolysis


**Step: 1,3-Bisphosphoglycerate → 3-Phosphoglycerate**

- **Enzyme:** Phosphoglycerate kinase
- **ATP Produced:** 1 per reaction
- **Reaction:**



**Step: Phosphoenolpyruvate (PEP) → Pyruvate**

- **Enzyme:** Pyruvate kinase
- **ATP Produced:** 1 per reaction
- **Reaction:**



## Significance of SLP

- Ensures ATP production during anaerobic conditions (e.g., intense exercise or low oxygen supply).
- Vital for red blood cells (which lack mitochondria and rely only on glycolysis).
- Important for quick energy bursts in muscle cells.
- Plays a key role in early evolution when oxidative phosphorylation was not yet developed.



## Inhibitors of Electron Transport Chain (ETC)

- These inhibitors block the transfer of electrons at specific points in the ETC, thereby halting ATP synthesis and oxygen consumption.

### 1. Complex I Inhibitors (NADH → Coenzyme Q)

- **Rotenone** (a plant-derived insecticide)
- **Amytal** (a barbiturate drug)

#### Mechanism:

- Inhibit **NADH dehydrogenase**
- Prevent the transfer of electrons from NADH to Coenzyme Q (ubiquinone)
- **FADH<sub>2</sub> can still donate electrons via Complex II**

#### Effect:

- ATP production from NADH is blocked
- ETC partially functions using FADH<sub>2</sub>

### 2. Complex II Inhibitors

- **Malonate** (competitive inhibitor of succinate dehydrogenase)

#### Mechanism:

- Inhibits **succinate dehydrogenase**, blocking electron entry from FADH<sub>2</sub>
- **Less severe effect**, since NADH can still enter via Complex I

### 3. Complex III Inhibitor (CoQ → Cytochrome c)

- **Antimycin A** (an antibiotic)

#### Mechanism:

- Inhibits the **cytochrome bc<sub>1</sub> complex**

- Prevents electron flow from reduced CoQ (ubiquinol) to cytochrome c

**Effect:**

- Entire chain is blocked beyond Complex III
- Both NADH and FADH<sub>2</sub> electron entry is ineffective

4. Complex IV Inhibitors (Cytochrome c → O<sub>2</sub>)


- **Cyanide (CN<sup>-</sup>)**
- **Carbon monoxide (CO)**
- **Sodium azide (NaN<sub>3</sub>)**

**Mechanism:**

- Inhibit **cytochrome c oxidase**
- Prevent the reduction of oxygen to water (final step of ETC)

**Effect:**

- Electron transport completely stops
- **Rapid ATP depletion**
- **Highly toxic and lethal**, especially cyanide



## Uncouplers of Oxidative Phosphorylation

→ Uncouplers disrupt the link between electron transport and ATP synthesis by dissipating the proton gradient. They make the inner mitochondrial membrane permeable to protons ( $H^+$ ), bypassing ATP synthase.

### 1. 2,4-Dinitrophenol (DNP)

- Synthetic chemical used historically for weight loss (now banned)

#### Mechanism:

- Acts as a **protonophore**: transports protons across the inner mitochondrial membrane
- Collapses the **proton gradient**, so **ATP synthase cannot function**

#### Effect:

- ETC continues to function and consumes oxygen
- **No ATP formed**
- Energy is **released as heat** → risk of **hyperthermia and death**

### 2. Thermogenin (UCP<sub>1</sub> - Uncoupling Protein 1)

- Naturally found in **brown adipose tissue** of infants and hibernating animals

#### Mechanism:

- Forms a channel for protons to re-enter the mitochondrial matrix
- **Bypasses ATP synthase**

#### Effect:

- ATP not produced
- Heat is generated instead (used for **thermoregulation**)

### 3. Aspirin (in high doses)

- High concentrations of **salicylate** act as a mild uncoupler

#### **Mechanism:**

- Increases mitochondrial membrane permeability to protons

#### **Effect:**

- Leads to heat production and **fever**
- Explains **fever in aspirin overdose**



**Pharmacy**  
Learn and Educate